

Topology Vol. 31, No. 4, pp. 801–804, 1992.
Printed in Great Britain

0040-9383/92 \$5.00 + 0.00
© 1992 Pergamon Press Ltd

ONE RELATOR GROUPS ARE SEMISTABLE AT INFINITY

MICHAEL L. MIHALIK and STEVEN T. TSCHANTZ

(Received 27 July 1990)

IN THIS paper, we show that all finitely generated one relator groups satisfy a homotopy property. For a connected CW-complex Y proper rays $r, s: [0, \infty) \rightarrow Y$ converge to the same end of Y if for any compact set $C \subseteq Y$ there is an integer N such that $r([N, \infty))$ and $s([N, \infty))$ are contained in the same component of $Y - C$. The space Y is *semistable at infinity* if any two proper rays converging to the same end of Y are properly homotopic. A finitely presented group G is *semistable at infinity* if for some (equivalently any) finite complex X with $\pi_1(X) = G$ the universal cover of X is semistable at infinity. Our main result is

THEOREM 1. *All finitely generated one relator groups are semistable at infinity.*

By [1] we have

COROLLARY 2. *For any one relator group G , $H^2(G; \mathbb{Z}G)$ is free Abelian.*

Our main tool in proving Theorem 1 is the following result from [3].

THEOREM 3. *If $G = A *_H B$ is an amalgamated product where A and B are finitely presented and semistable at infinity, and H is finitely generated then G is semistable at infinity. If $G = A *_H$ is an HNN-extension where A is finitely presented and semistable at infinity and H is finitely generated then G is semistable at infinity.*

Proof of Theorem 1. We use the following structure theorem for one relator groups patterned after Magnus' proof of the Freiheitssatz (see [2]).

LEMMA 4. *Given any finitely generated one relator group G there exists a finite sequence of finitely generated one relator groups $H_1, H_2, \dots, H_n = G$ such that for each $i < n$, either H_{i+1} or $H_{i+1} * \mathbb{Z}$ is an HNN-extension of H_i over a finitely generated group, and such that H_1 is either a free group or else is a free product of a free group and a finite cyclic group.*

The proof of the theorem also uses the following fact.

LEMMA 5. *If G is finitely presented and $G * \mathbb{Z}$ is semistable at infinity then G is semistable at infinity.*

Then each H_i is seen to be semistable at infinity by induction on i . Since free groups and finite groups are semistable at infinity, H_1 is semistable at infinity. If H_i is semistable at infinity then, by Theorem 3, an HNN-extension of H_i over a finitely generated group is semistable at infinity, so either H_{i+1} or $H_{i+1} * \mathbb{Z}$ is semistable at infinity. In the later case, by

Lemma 5, H_{i+1} is semistable at infinity again. But then by induction, each H_i is semistable at infinity and so $G = H_n$ is semistable at infinity. \blacksquare

Proof of Lemma 4. The proof is by induction on the length N of a cyclically reduced relator r defining G and for each fixed N by induction on $N - k$ where k is the number of generators appearing in r . If $N = 0$ then G is a free group. If $N > 0$ and $N - k = 0$, i.e. each generator in r appears exactly once, then G is a free group with one fewer generator. Otherwise, it will suffice to find a finitely generated one relator group H with defining relator r' of length N' involving k' generators such that either $N' < N$ or else $N' = N$ and $N' - k' < N - k$, and such that either G or $G * \mathbb{Z}$ is an HNN-extension of H over a finitely generated group.

First suppose that some letter a appearing in r has zero exponent sum in r . For each other generator b appearing in r define $b_i = a^{-i}ba^i$ and rewrite r as a product r' of the b_i (different b_i corresponding to each different b). Since there is at least one occurrence of a in r and each other b is rewritten as a single b_i , r' is shorter than r . For each b in r , let m_b and M_b be the greatest lower bound and least upper bound on the i with b_i appearing in r' . Let H have as generators the set of all b_i , $m_b \leq i \leq M_b$, together with any generators of G not appearing in r (so a is excluded) and single relator r' . Let K be the subgroup of H generated by all of the b_i , $m_b \leq i < M_b$. Then K is a free group freely generated by these b_i since the generators of K omit one of the generators of H appearing in r' . Similarly, let K' be the subgroup of H generated by all of the b_i , $m_b < i \leq M_b$, so that K' is also free and hence isomorphic to K by the isomorphism mapping b_i to b_{i+1} . But then the HNN-extension of H by these isomorphic subgroups with stable letter a has presentation $\langle a, b, \dots : r', a^{-1}b_i a = b_{i+1} (m_b \leq i < M_b) \rangle$. Defining additional b_i as needed, we can derive $b_i = a^{-i}b_0 a^i$ and so defining $b = b_0$ we get back the presentation of G , i.e. G is an HNN-extension of H and this case is complete.


Suppose now that no letter appears in r with zero exponent sum. If only one letter appears in r we have a group isomorphic to a free product of a free group and \mathbb{Z}_N . If more than one letter appears in r then there will be two cases depending on whether or not there are distinct letters a and b in r satisfying a certain condition. If such a and b exist then we construct a one relator group H with relator of the same length as r but involving more generators, such that $G * \mathbb{Z}$ is an HNN-extension of H over a finitely generated group. The condition on a and b will be devised just so this construction works. If no a and b can be found satisfying the condition then we show that G is a free product of a free group and a finite cyclic group.

Let x be a letter distinct from the generators of G so $G' = G * \mathbb{Z}$ has presentation $\langle a, b, \dots, x : r \rangle$. Let a and b be distinct letters appearing in r and let $\alpha = \sigma_a(r)$ and $\beta = \sigma_b(r)$ (where as usual $\sigma_a(r)$ denotes the exponent sum of the letter a in r .) Add new generators to this presentation with defining relations $y = ax^\beta$ and $z = bx^{-\alpha}$. Solve for a and b in terms of y and z , substitute $a = yx^{-\beta}$ and $b = zx^\alpha$ back into r to get a word r_1 , and eliminate the now redundant a and b . Hence G' has presentation $\langle y, z, \dots, x : r_1 \rangle$ where y and z have replaced a and b in the list of generators and where now $\sigma_x(r_1) = \sigma_a(r)(-\beta) + \sigma_b(r)\alpha = 0$. Proceed as in the previous case to rewrite r_1 as a word r'_1 in $c_i = x^{-i}cx^i$ where c ranges over the letters appearing in r_1 except x . Now r'_1 will have the same length as r since each a becomes a y_i , each b becomes a z_i and each other c becomes a c_i in r'_1 . Taking $H = \langle y_i, \dots, z_i, \dots : r'_1 \rangle$ we get that $G' = G * \mathbb{Z}$ is an HNN-extension of H over a finitely generated (free) group with stable letter x . If r'_1 involves more generators than r then we take $r' = r'_1$ and we are done (since $N' = N$ and $N' - k' < N - k$). We must examine this rewriting procedure in more detail to see exactly when this is the case.

Each occurrence of a letter c in r corresponds to a triple (s, u, t) where $u = c^{\pm 1}$, and $r = sut$. Define $I_{a,b}(s, u, t)$ for distinct letters a and b , $u = c^{\pm 1}$ for a letter c , and words s and t as follows. If $u \neq a^{-1}, b^{-1}$, let $I_{a,b}(s, u, t) = \sigma_a(s)\sigma_b(sut) - \sigma_b(s)\sigma_a(sut)$. Let $I_{a,b}(s, a^{-1}, t) = (\sigma_a(s) - 1)\sigma_b(sut) - \sigma_b(s)\sigma_a(sut)$, and $I_{a,b}(s, b^{-1}, t) = \sigma_a(s)\sigma_b(sut) - (\sigma_b(s) - 1)\sigma_a(sut)$. If $r = sut$ and $u = c^{\pm 1}$ then $\sigma_a(sut) = \alpha$, $\sigma_b(sut) = \beta$, and $I_{a,b}(s, u, t)$ is the negative of the exponent sum of x in the initial segment of r_1 before the corresponding occurrence of u in r_1 (or if $u = a^{-1}$ or b^{-1} , the negative of the exponent sum of x in the initial segment of r_1 before the corresponding y^{-1} or z^{-1}), and so this occurrence of $c^{\pm 1}$ in r becomes a $c_i^{\pm 1}$ in r'_1 where $i = I_{a,b}(s, u, t)$ (or $y_i^{\pm 1}$ or $z_i^{\pm 1}$ if $u = a^{\pm 1}$ or $b^{\pm 1}$). Thus more generators appear in r'_1 than appear in r provided that for some occurrences of a letter c in $r = sut = s'u't'$, $u = c^{\pm 1}$ and $u' = c^{\pm 1}$, we have $I_{a,b}(s, u, t) \neq I_{a,b}(s', u', t')$.

Now it may happen that for a particular a and b , r'_1 turns out to be just a relabeling of r and yet for a different choice of a and b , r'_1 would involve more generators than r and we can proceed as outlined above. Suppose instead that for every pair of distinct generators a and b appearing in r and for every pair of occurrences of c in $r = sut = s'u't'$, $u = c^{\pm 1}$ and $u' = c^{\pm 1}$ we have $I_{a,b}(s, u, t) = I_{a,b}(s', u', t')$. We now show that in this case G is a free product of a free group and a finite cyclic group.

First we may assume that for each letter c occurring in r , $\sigma_c(r) > 0$. Suppose that for some a and b , $\sigma_a(r) > \sigma_b(r)$ and take $\alpha = \sigma_a(r)$ and $\beta = \sigma_b(r)$ as usual. Partition r as a product $s_0 a s_1 a s_2 \dots a s_n$ where each s_i has $\sigma_a(s_i) = 0$. By conjugating r we may as well assume that $s_0 = 1$. Then $\alpha > \beta = \sigma_b(r) = \sum_{i=1}^n \sigma_b(s_i)$ is a sum of α terms so some $\sigma_b(s_i) \leq 0$. By conjugating again we may as well assume that $\sigma_b(s_1) \leq 0$. But then $I_{a,b}(1, a, s_1 a \dots s_n) = 0$ and $I_{a,b}(a s_1, a, s_2 \dots s_n) = \beta - \sigma_b(a s_1) \alpha > 0$ and so this choice of a and b would be suitable for the above construction. Hence instead $\sigma_a(r)$ is the same for all a , say $\sigma_a(r) = m$. Next suppose that for some c both c and c^{-1} appear in r (still assumed to be cyclically reduced). Take a to be a letter such that $r = s_0 a s_1 a^{-1} s_2$ or else $r = s_0 a^{-1} s_1 a s_2$ with s_1 of minimal length (over all possible pairs of occurrences of letters and inverses). Say the former case holds and let b be a letter occurring in s_1 . Then $\sigma_a(s_1) = 0$ since there can be no occurrence of a or a^{-1} in s_1 , and $\sigma_b(s_1) \neq 0$ since not both of b and b^{-1} appear in s_1 by the minimality assumption. But then $I_{a,b}(s_0 a s_1, a^{-1}, s_2) = I_{a,b}(s_0, a, s_1 a^{-1} s_2) - \sigma_b(s_1) m$ and again a and b would work in the above construction.



Hence each letter appears the same number of times with only positive exponents. If there exists a and b such that b does not occur or appears more than once between some pair of successive occurrences of a then these occurrences of a would become different y_i in the above construction. Hence if there are k generators and a is the first letter in r then each other generator must appear exactly once between the first and second occurrences of a in r , i.e. the second occurrence of a is the $k + 1$ -st letter of r . But then the second occurrence of the second letter of r must be the $k + 2$ -nd letter of r , etc. The relator r must be a power of the product of the generators appearing in r , and hence G is a free product of a free group (generated by the generators of G not appearing in r) and a group with presentation $\langle a_1, \dots, a_k : (a_1 \dots a_k)^m \rangle$. But this last group is a free product of a free group of rank $k - 1$ and \mathbb{Z}_m . Thus, in the case where no letter appears in r with zero exponent sum, either there exist letters a and b such that the above construction produces a one relator group H with relator r' of the same length but involving more generators such that $G * \mathbb{Z}$ is an HNN-extension of H , or else the relator r is of such a simple form that G is in fact a free product of a free group and a finite cyclic group. ■

Proof of Lemma 5. Let X be a finite complex such that $\pi_1(X) = G$. Let Y be obtained from X by attaching a loop to X at a vertex $x \in X$. Then $\pi_1(Y) = G * \mathbb{Z}$. The universal cover \tilde{Y} of Y is a union of copies of \tilde{X} , the universal cover of X , and copies of the real line \mathbb{R} . Let $p: \tilde{Y} \rightarrow Y$ be the covering projection. If a copy of \tilde{X} intersects a copy of \mathbb{R} it does so in exactly one point of $p^{-1}(x)$. If a copy of \tilde{X} intersects a copy of \mathbb{R} it does so in exactly one point of $p^{-1}(x)$. All copies of \tilde{X} are mutually disjoint as are all copies of \mathbb{R} (see figure).

Let \tilde{X}_0 be a copy of \tilde{X} in \tilde{Y} . Let $r, s: [0, \infty) \rightarrow \tilde{X}_0$ be proper rays converging to the same end of \tilde{X}_0 . It suffices to show that r and s are properly homotopic in \tilde{X}_0 . Since \tilde{Y} is semistable at infinity there exists $F: [0, \infty) \times [0, 1] \rightarrow \tilde{Y}$ a proper homotopy of r to s . If U is a component of $\tilde{Y} - \tilde{X}_0$ then \bar{U} , the closure of U in \tilde{Y} , is $U \cup \{v\}$ for some vertex $v \in p^{-1}(x)$. Since no proper ray in \bar{U} converges to the same end of \tilde{Y} as a ray in \tilde{X}_0 , $F^{-1}(\bar{U})$ is compact. If a is any point in the topological boundary of $F^{-1}(\bar{U})$ then $F(a) \in \bar{U} - U = \{v\}$. Define $F': [0, \infty) \times [0, 1] \rightarrow \tilde{X}_0$ as follows. For $a \in F^{-1}(\tilde{X}_0)$ let $F'(a) = F(a)$. For $a \in F^{-1}(U)$ where U is a component of $\tilde{Y} - \tilde{X}_0$ and v the unique point of $\bar{U} - U$ let $F'(a) = v$. If C is a compact subset of \tilde{X}_0 let U_1, \dots, U_n be the components of $\tilde{Y} - \tilde{X}_0$ whose closures intersect C (finitely many since $C \cap p^{-1}(x)$ is finite). Then $(F')^{-1}(C) = F^{-1}(C) \cup \bigcup_{i=1}^n F^{-1}(\bar{U}_i)$. It follows that F' is proper and continuous. ■

REFERENCES

1. R. GEOGHEGAN and M. MIHALIK: Free abelian cohomology of groups and ends of universal covers, *J. Pure and Appl. Algebra* **36** (1985), pp 123–137.
2. R. C. LYNDON and P. E. SCHUPP: *Combinatorial Group Theory*, Springer-Verlag, Berlin, Heidelberg, New York (1970).
3. M. MIHALIK and S. TSCHANTZ: Semistability of amalgamated products and HNN-extensions, *Mem. Am. Math. Soc.*, to appear.

*Department of Mathematics,
Vanderbilt University,
Nashville,
Tennessee 37240,
U.S.A.*